arXiv:1208.3021v1 [gr-qc] 15 Aug 2012

Separation of spacetime and matter in polar oscillations of compact stars

Y. J. Zhang*, Q. Q. Zhu, J. Wu†, and P. T. Leung‡
Physics Department and Institute of Theoretical Physics,
The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China.
(Dated: August 16, 2012)

In polar oscillations of compact stars dynamical changes in spacetime and matter are in general coupled together. Based on the theory of stellar pulsation proposed by Allen *et al.* [Phys. Rev. D **58**, 124012 (1998)], we establish a feasible approximation scheme to separate spacetime from matter in such oscillations. Two independent second-order ordinary differential equations are obtained, which lead to accurate determination of p-mode and w-mode quasi-normal modes of compact stars, respectively.

PACS numbers: 04.40.Dg, 04.30.Db, 97.60.Jd, 95.30.Sf

I. INTRODUCTION

Compact stars, including neutron stars (NSs) and quark stars (QSs), are notable for the high density achievable at their centers and the huge gravitational field surrounding them. The static structure of NSs (or QSs) is often the ideal test bed for the equation of state (EOS) of nuclear (or quark) matter and can provide clues to the interactions between elementary particles as well (see, e.g., [1–3] and references therein). Likewise, gravitational waves (GWs) emitted from oscillating NSs (or QSs) are also expected to carry rich information about the internal structure of these stars, giving rise to various efforts to infer the properties of relativistic compact stars from close examination of their oscillation spectra (see, e.g., [4–12]). Although the oscillation frequencies of NSs usually lie outside the sensitivity window of the existing GW detectors such as GEO600, LIGO I, TAMA300 and VIRGO (see, e.g., [13, 14]), with the advent of more sensitive GW telescopes (e.g., the proposed Einstein Telescope [15, 16]), it might be possible to infer the structure and the composition of a compact star from its GW signals in the coming future.

The fully relativistic formulation of stellar pulsations was founded by Thorne and Campolattaro in 1967 [17] and since then there have been other attempts to describe such processes with other alternative approaches (see, e.g., [18–22]). Basically, oscillations of compact stars are classified into axial and polar types. For axial type oscillations, the matter forming a star is essentially a spectator and only the spacetime surrounding the star pulsates in accordance with a scalar wave equation [20]. On the other hand, in polar type oscillations matter motion and spacetime variation are coupled together and hence polar type oscillations are expected to be more relevant to dynamical processes such as supernova and binary mergers of NSs.

To describe the interplay between matter and spacetime in polar oscillations of compact stars, different versions of equations of motion governing such coupled motion have been proposed since the pioneering work of Thorne and Campolattaro [17]. For example, based on [17], Lindblom and Detweiler described the system with four coupled first-order differential equations (termed as the LD formalism in this paper) [18, 19], which can be readily solved numerically. Later, to draw an analogy between the oscillations of relativistic and Newtonian stars, Lindblom, Mendell and Ipser [21] proposed a new formalism to describe the coupled oscillation with two second-order equations (termed as the LMI formalism in this paper), which reduce directly into the corresponding Newtonian equations under the weak field limit.

In another relativistic stellar pulsation formalism proposed by Allen, Anderson, Kokkotas and Schutz (termed as the AAKS formalism in this paper) [23], variations in matter and spacetime are described by one matter variable and two metric variables, which are governed by three coupled second-order time-dependent wave equations. In fact, these three variables are also related through a time-independent Hamiltonian constraint. Hence, only two of the three wave equations are independent. Allen *et al.* [23] managed to evolve these variables in the time domain.

The aim of the present paper is to study polar pulsations of compact stars with the AAKS formalism in the frequency domain where all dynamical variables are assumed to have $\exp(-i\omega t)$ time dependence, with ω being the eigenfrequency. Oscillation modes in which GWs are emitted from a star to spatial infinity are termed as quasi-normal modes (QNMs) because they are usually damped [24–28]. Hence, the eigenfrequency $\omega \equiv \omega_{\mathrm{r}} - i\omega_{\mathrm{i}}$ is complex-valued with $\omega_{\mathrm{i}} \geq 0$ measuring the decay rate. QNMs of compact stars are worth studying because, as mentioned above, they

---

* Present address: Department of Physics, University of California at San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA
† Present address: Physics Department, Hong Kong Baptist University, 224 Waterloo Road, Kowloon Tong, Hong Kong SAR, China.
‡ Email: ptleung@phy.cuhk.edu.hk

can reveal the physical characteristics (e.g., mass, radius, moment of inertia, equation of state (EOS) and composition) of a star (see, e.g., [4–11]). On the other hand, polar QNMs of compact stars can be classified into the fluid mode (including the fundamental f-mode, the pressure p-mode and the gravity g-mode according to the nature of respective restoring forces [29]) and the spacetime w-mode which has no Newtonian counterpart.

In the present paper, based on the AAKS formalism, we derive in the frequency domain two coupled wave equations governing the evolution of the matter field and the metric variables. As in the LD formalism, these two equations together lead to both fluid and spacetime QNMs of compact stars. In addition, we show that these two equations can be straightforwardly decoupled, resulting in two independent equations which can reproduce good approximation of fluid modes and w-modes, respectively. With such decoupling scheme for matter and spacetime, we succeed in obtaining the Cowling approximation (CA), where effects of metric perturbations are neglected [21, 30], and the inverse-Cowling approximation (ICA), where fluid motion is ignored [31, 32], for compact stars in a unified framework. The fluid motion is described by a single second-order equation in the matter field without involving any spacetime perturbations. Meanwhile, polar w-mode oscillations obey a single wave equation as in the case of axial w-modes [20]. To our knowledge, this is the first time a single second-order equation for polar w-mode QNMs is found.

Furthermore, we show that the decoupled equation for fluid motion, upon imposing suitable physical boundary conditions, form a Sturm-Liouville eigenvalue problem (see, e.g., [33]). Therefore, the fluid system can be studied with conventional normal-mode analysis, and the energy of the fluid is conserved as long as the influence of the spacetime perturbations are negligible. The mode with the lowest eigenfrequency corresponds to the f-mode oscillation, while other modes are good approximation for p-mode QNMs.

The organization of the present paper is as follows. We briefly review the equilibrium state and the perturbed state of a compact star in Secs. II and III, respectively. The AAKS formalism for pulsation of compact stars in the frequency domain is developed in Sec. IV. In Secs. V and VI we recast the equations of motion in the AAKS formalism into two coupled wave equations in two variables. We show in Secs. VII and VIII that these two equations can be decoupled, respectively leading to accurate CA and ICA schemes. We conclude the present paper with a discussion in Sec. IX. Besides, there are two appendices respectively providing detailed discussion on the boundary condition imposed at the stellar surface and an alternative approach to description of pulsation of compact stars.

Geometric units in which $G = c = 1$ are adopted throughout this paper. We consider polar oscillations of a non-rotating compact star with radius R and total mass M. Unless stated otherwise, all numerical values of frequencies are evaluated for the case where the angular momentum index l equals 2 and measured in units of M^{-1}. Besides, we also assume that the effect of temperature on the EOS of nuclear matter is negligible and hence g-modes are omitted in our discussion.

II. EQUILIBRIUM STATE OF COMPACT STARS

The energy-momentum tensor of a non-rotating compact star made of a perfect fluid at equilibrium state is given by

$$T^{\alpha\beta} = (p+\rho)u^\alpha u^\beta + p g^{\alpha\beta}, \tag{2.1}$$

where p is the pressure, ρ is the energy density and u^α is the 4-velocity of the fluid. The spacetime around the star is described by the line element:

$$ds^2 = g_{\alpha\beta}dx^\alpha dx^\beta = -e^\nu dt^2 + e^\lambda dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2), \tag{2.2}$$

where $(x^0, x^1, x^2, x^3) = (t, r, \theta, \phi)$, and reduces to the standard Schwarzschild form outside the star,

$$ds^2 = -(1-\frac{2M}{r})dt^2 + (1-\frac{2M}{r})^{-1}dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2). \tag{2.3}$$

By substituting $T^{\alpha\beta}$ and $g_{\alpha\beta}$ into the Einstein equation, the Tolman-Oppenheimer-Volkov (TOV) equations of hydrostatic equilibrium can be derived [34, 35]:

$$\begin{aligned} \frac{dp}{dr} &= -\frac{(\rho+p)(m+4\pi r^3 p)}{r(r-2m)}, && (2.4) \\ \frac{d\nu}{dr} &= \frac{2(m+4\pi r^3 p)}{r(r-2m)}, && (2.5) \\ e^{-\lambda} &= 1-2m/r, && (2.6) \end{aligned}$$

with

$$m(r) = \int_0^r 4\pi r'^2 \rho(r') dr'. \tag{2.7}$$

The hydrostatic structure of a compact star can be specified by the TOV equations, together with the star's EOS $p = p(\rho)$ and the boundary conditions:

$$\begin{aligned} e^{\nu}|_{r=R} &= 1 - 2M/R, &(2.8)\\ p|_{r=R} &= 0. &(2.9)\end{aligned}$$

III. PERTURBED NEUTRON STARS

When a static compact star is perturbed, its fluid and spacetime perform small-amplitude oscillations about the equilibrium state. Hence, the perturbed metric is given by:

$$\widetilde{g}_{\alpha\beta} = g_{\alpha\beta} + h_{\alpha\beta}, \tag{3.1}$$

where $g_{\alpha\beta}$ is the spacetime metric of the equilibrium configuration given by the TOV equations [34, 35]. The perturbations in the spacetime metric and fluid displacement are first decomposed into tensorial and vectorial spherical harmonics with angular momenta $\{lm\}$, and then separated into axial and polar parts according to parity. The polar part of the perturbation in the metric, h^{p}_{lm}, in the Regge-Wheeler gauge can be written as [17, 36, 37]:

$$(h^{\mathrm{p}}_{lm})_{\alpha\beta} = \begin{pmatrix} e^{\nu}\bar{H}_0 & \bar{H}_1 & 0 & 0 \\ \bar{H}_1 & e^{\lambda}\bar{H}_0 & 0 & 0 \\ 0 & 0 & r^2\bar{K} & 0 \\ 0 & 0 & 0 & r^2\sin^2\theta\bar{K} \end{pmatrix} Y_{lm}, \tag{3.2}$$

and the corresponding fluid displacements are [17–19]:

$$\begin{aligned} \xi^r &= r^{-1}e^{-\lambda/2}\bar{W}Y_{lm}, &(3.3)\\ \xi^{\theta} &= -r^{-2}\bar{V}\partial_{\theta}Y_{lm}, &(3.4)\\ \xi^{\phi} &= -(r\sin\theta)^{-2}\bar{V}\partial_{\phi}Y_{lm}, &(3.5)\end{aligned}$$

where $\bar{H}_0$, $\bar{H}_1$, $\bar{K}$, $\bar{W}$ and $\bar{V}$ are functions of both t and r.

IV. AAKS FORMALISM IN FREQUENCY DOMAIN

A. Equations inside the Star

In the AAKS formalism [23], polar oscillations of compact stars are described by two spacetime metric variables $\bar{F}(t,r)$, $\bar{S}(t,r)$ and one fluid perturbation variable $\bar{H}(t,r)$ which are related to the metric and the Eulerian change in pressure, δp, as follows:

$$\begin{aligned} \bar{S}(r,t) &= \frac{e^{\nu}\left[\bar{H}_0(r,t) - \bar{K}(r,t)\right]}{r}, &(4.1)\\ \bar{F}(r,t) &= r\bar{K}(r,t), &(4.2)\\ \bar{H}(r,t) &= \frac{\delta p(r,t)}{\rho + p}. &(4.3)\end{aligned}$$

For QNMs, the three variables $\bar{S}(r,t)$, $\bar{F}(r,t)$ and $\bar{H}(r,t)$ all have the $e^{-i\omega t}$ time-dependence, i.e.,

$$\begin{aligned} \bar{S}(r,t) &= S(r)e^{-i\omega t}, &(4.4)\\ \bar{F}(r,t) &= F(r)e^{-i\omega t}, &(4.5)\\ \bar{H}(r,t) &= H(r)e^{-i\omega t}. &(4.6)\end{aligned}$$

It is readily shown that $S(r)$, $F(r)$ and $H(r)$ satisfy the following three coupled second-order differential equations [23]:

$$\begin{aligned}
&\omega^2 S+\frac{d^2S}{dr_*^2}+\frac{2e^\nu}{r^3}\left[2\pi r^3(\rho+3p)+m-(n+1)r\right]S \\
=\;& -\frac{4e^{2\nu}}{r^5}\left[\frac{(m+4\pi pr^3)^2}{r-2m}+4\pi\rho r^3-3m\right]F, \qquad (4.7)\\
&\omega^2 F+\frac{d^2F}{dr_*^2}+\frac{2e^\nu}{r^3}\left[2\pi r^3(3\rho+p)+m-(n+1)r\right]F \\
=\;& -2\left[4\pi r^2(p+\rho)-e^{-\lambda}\right]S+8\pi(\rho+p)re^\nu(1-\frac{1}{C_s^2})H, \qquad (4.8)\\
&\frac{\omega^2}{C_s^2}H+\frac{d^2H}{dr_*^2}+\frac{e^{(\nu+\lambda)/2}}{r^2}\frac{dH}{dr_*}\left[(m+4\pi pr^3)(1-\frac{1}{C_s^2})+2(r-2m)\right]\\
&+\frac{2e^\nu}{r^2}\left[2\pi r^2(\rho+p)(3+\frac{1}{C_s^2})-(n+1)\right]H\\
=\;&\left[\frac{(m+4\pi pr^3)^2}{r^2(r-2m)}(1+\frac{1}{C_s^2})-\frac{m+4\pi pr^3}{2r^2}(1-\frac{1}{C_s^2})-4\pi r(3p+\rho)\right]S\\
&+\frac{e^\nu}{r^2}\left[\frac{2(m+4\pi pr^3)^2}{r^2(r-2m)}\frac{1}{C_s^2}-\frac{m+4\pi pr^3}{2r^2}(1-\frac{1}{C_s^2})-4\pi r(3p+\rho)\right]F\\
&+(m+4\pi pr^3)(1-\frac{1}{C_s^2})\frac{e^{(\lambda-\nu)/2}}{2r}\left(\frac{e^\nu}{r^2}\frac{dF}{dr_*}-\frac{dS}{dr_*}\right), \qquad (4.9)
\end{aligned}$$

where

$$n=(l+2)(l-1)/2, \qquad (4.10)$$

r_* is the standard tortoise coordinate related to the circumferential radius r through the relationship:

$$r_*(r)=\int_0^r \exp\left[\frac{\lambda(r')-\nu(r')}{2}\right]dr', \qquad (4.11)$$

and $C_s^2(r)$ is the square of the sound speed in the stellar fluid, defined as

$$C_s^2=dp/d\rho. \qquad (4.12)$$

In addition to the above three equations, $S(r)$, $F(r)$ and $H(r)$ should also satisfy the following Hamiltonian constraint:

$$\begin{aligned}
&\frac{d^2F}{dr_*^2}-\frac{e^{(\nu+\lambda)/2}}{r^2}(m+4\pi r^3p)\frac{dF}{dr_*}+\frac{e^\nu}{r^3}[12\pi r^3\rho-m-2(n+1)r]F\\
-\;& re^{(\nu+\lambda)/2}\frac{dS}{dr_*}+\left[8\pi r^2(p+\rho)-(n+3)+\frac{4m}{r}\right]S+\frac{8\pi r}{C_s^2}e^\nu(\rho+p)H=0. \qquad (4.13)
\end{aligned}$$

By combining (4.8) and the Hamiltonian constraint (4.13) together, a simple expression for H can be derived [23]:

$$8\pi(\rho+p)re^\nu H \;=\; \frac{e^{(\lambda+\nu)/2}}{r^2}(m+4\pi pr^3)\frac{dF}{dr_*}+\omega^2F+\frac{e^\nu}{r^3}(3m+4\pi pr^3)F+e^{-(\nu+\lambda)/2}r\frac{dS}{dr_*}+(n+1)S, \qquad (4.14)$$

which is often used to calculate the value of H, instead of integrating (4.9), and termed as the modified Hamiltonian constraint hereafter. However, Eq. (4.14) obviously fails to give the value of H at the stellar surface, where both p and ρ vanish. We will further discuss such subtlety in the later part of this section.

B. Equations outside the Star

As for the region outside the star the fluid variable H vanishes identically, the two metric variables S and F evolve as:

$$\omega^2 S + \frac{d^2S}{dr_*^2} + \frac{2(r-2M)}{r^4}\left[M-(n+1)r\right]S = -\frac{4M}{r^7}(r-2M)(7M-3r)F, \tag{4.15}$$

$$\omega^2 F + \frac{d^2F}{dr_*^2} + \frac{2(r-2M)}{r^4}\left[M-(n+1)r\right]F = 2(1-\frac{2M}{r})S, \tag{4.16}$$

which can be obtained from (4.7) and (4.8) by setting $p=\rho=0$ and $m(r)=M$. In addition, they are related to the Zerilli function $Z(r_*)$ as follows:

$$\frac{(n+1)(nr+3M)}{r^3}Z = -\frac{1}{r}\frac{dF}{dr_*} + \frac{(n+2)r-M}{r^3}F + S, \tag{4.17}$$

where the Zerilli function $Z(r_*)$ satisfies the Zerilli equation [38]:

$$\left[\frac{d^2}{dr_*^2} + \omega^2 - V_Z(r_*)\right]Z(r_*) = 0, \tag{4.18}$$

with the Zerilli potential $V_Z(r_*)$ given by

$$V_Z(r_*) = \frac{r-2M}{r^4(nr+3M)^2}\left[2n^2(n+1)r^3 + 6n^2Mr^2 + 18nM^2r + 18M^3\right]. \tag{4.19}$$

C. Evaluation of quasi-normal modes

QNMs of vibrating compact stars are defined by the physical solutions of the AAKS equations with the Zerilli function satisfying the outgoing boundary condition $Z \to \exp(i\omega r_*)$ at spatial infinity. In solving the AAKS equation set inside the star, we numerically integrate (4.7), (4.8) outward to find S and F, and use the modified Hamiltonian constraint (4.14) to evaluate H. However, a few remarks are noteworthy in carrying out the numerical integration scheme from $r=0$ to $r=R$.

First of all, there is a regular singular point at $r=0$ in (4.7)-(4.9) resulting from the term $(n+1)/r^2$, which forbids normal discretization scheme. Instead, the regularity boundary condition for S, F and H at $r=0$ leads to the following asymptotic behavior near the origin:

$$\begin{aligned} S &\approx s_0 r^{l+1}, & (4.20)\\ F &\approx f_0 r^{l+1}, & (4.21)\\ H &\approx h_0 r^{l}, & (4.22)\end{aligned}$$

where f_0, s_0 and h_0 are constants. Expanding the modified Hamiltonian constraint (4.14) about the origin, we find that these three constants satisfy the following relationship:

$$2(\rho_0+p_0)e^{\nu_0}h_0 = [e^{\nu_0/2}(\rho_0/3+p_0)(l+1)+\omega^2+e^{\nu_0}(\rho_0+p_0)]f_0 + [e^{-\nu_0/2}(l+1)+n+1]s_0, \tag{4.23}$$

where ρ_0, p_0 and $e^{-\nu_0}$ stand for the values of ρ, p and $e^{-\nu}$ evaluated at $r=0$, respectively. We make use of the asymptotic behavior in (4.20), (4.21) and (4.22) and start numerical integration at a point slightly away from the point $r=0$. Owing to the relationship (4.23), two independent solutions can be obtained when the AAKS equations are integrated outward. We then impose a proper boundary condition at the stellar surface $r=R$ to fix the ratio of these two independent solutions.

Since the Lagrangian perturbation of pressure Δp vanishes at the stellar surface, we have

$$\delta p(r=R) = -\left[\xi^r\frac{dp}{dr}\right]_{r=R}. \tag{4.24}$$

The value of H there then follows directly from the definition of H, (4.3), and the TOV equations [39]:

$$H(r=R) = -\frac{\xi^r(R)M}{R(R-2M)}. \tag{4.25}$$

Together with the modified Hamiltonian constraint (4.14) and the fact that $\rho = p = 0$ at the stellar surface, the boundedness of $H(r=R)$ implied by the above equation leads to the following boundary condition at $r = R$:

$$\omega^2 F + \frac{M}{R^2}\frac{\partial F}{\partial r_*} + R\frac{\partial S}{\partial r_*} + \frac{3M}{R^3}(1-\frac{2M}{R})F + (n+1)S = 0. \tag{4.26}$$

However, we find that this is merely a necessary condition, instead of a sufficient one, and cannot be considered as an independent boundary condition.

Instead, another independent boundary condition can be obtained by keeping only terms of order $1/C_s^2$ in (4.9) as $C_s \to 0$ when approaching the stellar surface [39, 40]:

$$-\omega^2 H + \frac{M}{R^2}\frac{\partial H}{\partial r_*} - \frac{M}{2R^3}\frac{\partial F}{\partial r_*} + \frac{M}{2(R-2M)}\frac{\partial S}{\partial r_*} + \frac{M(R+2M)}{2R^5}F + \frac{M}{2R(R-2M)}S = 0. \tag{4.27}$$

By imposing this condition on the linear combination of the two independent solutions, we obtain a single (up to a multiplicative constant) solution of S, F and H inside the star, which is connected to the Zerilli function outside the star to determine the QNM frequencies.

However, a remark about the accuracy of the direct integration scheme mentioned above is in order. Since the modified Hamiltonian constraint (4.14) is used to find H, considerable numerical errors could arise near the stellar surface where both ρ and p are small. To overcome this problem, we analytically expand S, F and H as power series there and connect such expansion with the solution obtained from numerical integration starting from the center of the star at a point near the stellar surface (see Appendix A). We find that such a matching scheme can yield more accurate results than the direct integration scheme. Moreover, the expansion could shed light on the relationships among the three physical fields (S, F and H) near the stellar surface, which are useful to the formulation of the decoupling schemes developed in the later part of the present paper.

V. *H-S* APPROACH

As mentioned in the Introduction, polar QNMs can be divided into two categories, namely, fluid modes and spacetime modes. In fluid modes, especially the p-modes, the associated GW emission is almost negligible, while in the spacetime w-modes, the fluid is hardly involved. It strongly suggests that appropriate schemes can be developed to decouple spacetime oscillations and fluid motions, leading to reduction in the order and hence simplification of the system.

It has been shown that inside the star and in the weak field limit K and H_0 are related to the perturbation in the Newtonian potential, U_1, by [41]:

$$K \simeq H_0 \simeq -2U_1. \tag{5.1}$$

Therefore, following directly from (4.1) and (4.2), to leading orders in field strengths, S and F measure the relativistic and Newtonian effects of gravity, respectively. Hence, in order to describe generation of gravitational radiation, which is the dominant process in the spacetime mode, S is the right variable to keep. On the other hand, H is proportional to the Eulerian change in pressure and describes the motion of matter.

In the following, based on the AAKS formalism, we establish a two-equation approach and a decoupling scheme leading to accurate CA for the fluid modes (especially the p-modes) and ICA for the w-modes using H and S, which is termed as the H-S approach in the present paper. First of all, we express F in terms of S and H for the regions $r < R$ and $r > R$.

For the case $r < R$ we rewrite (4.14) as:

$$8\pi(\rho+p)e^{\nu}rH = A_{f1}\frac{dF}{dr_*} + A_{f0}F + A_{s1}\frac{dS}{dr_*} + A_{s0}S, \tag{5.2}$$

with

$$\begin{aligned}
A_{s1} &= e^{-(\nu+\lambda)/2} r, \qquad &(5.3)\\
A_{s0} &= n+1, &(5.4)\\
A_{f1} &= \frac{e^{(\lambda+\nu)/2}}{r^2}(m+4\pi p r^3), &(5.5)\\
A_{f0} &= \omega^2 + \frac{e^{\nu}}{r^3}(3m+4\pi p r^3). &(5.6)
\end{aligned}$$

Differentiating (5.2) once, and using (4.7) and (4.8) to eliminate d^2F/dr_*^2 and d^2S/dr_*^2, we get:

$$8\pi(\rho+p)e^{\nu}\frac{d}{dr_*}(rH) = B_{f1}\frac{dF}{dr_*} + B_{f0}F + B_{s1}\frac{dS}{dr_*} + B_{s0}S, \tag{5.7}$$

where

$$\begin{aligned}
B_{s1} &= 1+n+e^{-\lambda} - 4\pi r^2(p+\rho), &(5.8)\\
B_{s0} &= \frac{e^{(\nu-\lambda)/2}}{r}\left[2(1+n) - e^{-\nu}r^2\omega^2 - 4\pi r^2(p+\rho)\left(1+r\nu'\right)\right], &(5.9)\\
B_{f1} &= \omega^2 + e^{\nu}\left[\frac{m}{r^3} + 4\pi(2p+\rho)\right], &(5.10)\\
B_{f0} &= \frac{e^{(\nu-\lambda)/2}}{r^4}\left\{e^{\nu}(3m-4\pi r^3\rho) - \frac{r^2\nu'}{2}\left[r^2\omega^2 - 2e^{\nu}(1+n-8\pi r^2(p+\rho))\right]\right\}, &(5.11)
\end{aligned}$$

with $\nu' = d\nu/dr$.

From (5.2) and (5.7), we can express F and dF/dr_* in terms of S, dS/dr_*, H and dH/dr_*:

$$\begin{aligned}
F &= \alpha_{s1}\frac{dS}{dr_*} + \alpha_{s0}S + \alpha_{h1}\frac{dH}{dr_*} + \alpha_{h0}H, &(5.12)\\
\frac{dF}{dr_*} &= \beta_{s1}\frac{dS}{dr_*} + \beta_{s0}S + \beta_{h1}\frac{dH}{dr_*} + \beta_{h0}H, &(5.13)
\end{aligned}$$

where

$$\begin{aligned}
\alpha_{s1} &= -\frac{1}{D}\Bigg\{e^{-(\lambda+\nu)/2}r\omega^2 + \frac{1}{r^2}e^{(\nu-\lambda)/2}\left[4\pi r^3(p+\rho) - e^{\lambda}(m+4\pi r^3 p)(1+n)\right.\\
&\qquad \left. +4\pi r^2 e^{\lambda}(m+4\pi r^3 p)(p+\rho)\right]\Bigg\}, &(5.14)\\
\alpha_{s0} &= -\frac{1}{D}\Bigg\{\left[n+1+\frac{m}{r}+4\pi r^2 p\right]\omega^2 + \frac{1}{r^3}e^{\nu}\left[-(1+n)(m-4\pi r^3\rho)\right.\\
&\qquad \left. +4\pi r^2 e^{\lambda}(m+4\pi r^3 p)(1+8\pi r^2 p)(p+\rho)\right]\Bigg\}, &(5.15)\\
\alpha_{h1} &= -\frac{8\pi e^{(\lambda+3\nu)/2}(\rho+p)}{Dr}(m+4\pi p r^3), &(5.16)\\
\alpha_{h0} &= \frac{8\pi e^{\nu}(\rho+p)r}{D}\left[\omega^2 + 4\pi e^{\nu}(p+\rho)\right], &(5.17)
\end{aligned}$$

$$\begin{aligned}\beta_{s1} &= -\frac{1}{D}\Big\{ \Big[n+2-\frac{m}{r}-4\pi r^2\rho\Big]\omega^2 + \frac{1}{r^3}e^{\nu}\left\{m\left[1+n-4\pi r^2(p+\rho)\right]\right.\\ &\left. +4\pi r^3\left[\rho-np+12\pi r^2 p(p+\rho)\right]\right\}\Big\},\end{aligned}\tag{5.18}$$

$$\begin{aligned}\beta_{s0} &= -\frac{1}{D}e^{-(\lambda+\nu)/2}\Big\{ -r\omega^4+\frac{1}{r^2}e^{\nu}\left\{e^{\lambda}(m+4\pi r^3 p)\left[1+n-8\pi r^2(p+\rho)\right]\right.\\ &\left. +2r(n+1)-3m-4\pi r^3(2p+\rho)\right\}\omega^2+\frac{1}{r^4}e^{2\nu}\Big\{m(1+n+\frac{2m}{r})\\ &+2(m+4\pi r^3 p)\Big[\frac{m}{r}+n-e^{\lambda}(1+n)^2\Big]+4\pi r(\rho+p)(m+4\pi r^3 p)\\ &\left\{r-2e^{\lambda}\left[m-r(1+2n)+4\pi r^3 p\right]\right\}+4\pi r^3\rho[1+n-\frac{2m}{r}]\Big\}\Big\},\end{aligned}\tag{5.19}$$

$$\beta_{h1} = \frac{8\pi e^{\nu}(\rho+p)r}{D}\left[\omega^2+\frac{e^{\nu}}{r^3}(3m+4\pi p r^3)\right],\tag{5.20}$$

$$\begin{aligned}\beta_{h0} &= \frac{8\pi e^{(3\nu-\lambda)/2}(\rho+p)}{D}\Big\{\Big[1+\frac{1}{r}e^{\lambda}(m+4\pi r^3 p)\Big]\omega^2+e^{\nu}\frac{1}{r^3}\left\{4\pi r^3(p+\rho)\right.\\ &\left.-2e^{\lambda}(m+4\pi r^3 p)\left[1+n-8\pi r^2(p+\rho)\right]\right\}\Big\},\end{aligned}\tag{5.21}$$

and

$$\begin{aligned}D &= \frac{2e^{2\nu}}{r^6}\left\{8\pi r^3(m+2\pi r^3 p)(p+\rho)-e^{\lambda}(m+4\pi r^3 p)^2\left[1+n-8\pi r^2(p+\rho)\right]\right\}\\ &+\frac{e^{\nu}}{r^4}\left[e^{\lambda}(m+4\pi r^3 p)^2+4mr+4\pi r^4(3p+\rho)\right]\omega^2+\omega^4.\end{aligned}\tag{5.22}$$

For $r > R$, the stellar pressure p and density ρ vanish and so does H, then F in (5.12) reduces to the following form:

$$F = \alpha_{s1}^{(\mathrm{e})}\frac{dS}{dr_*}+\alpha_{s0}^{(\mathrm{e})}S,\tag{5.23}$$

which is a linear combination of S and dS/dr_*. The coefficients $\alpha_{s0}^{(\mathrm{e})}$ and $\alpha_{s1}^{(\mathrm{e})}$ are obtained from α_{s0} and α_{s1} by taking $\rho = p = 0$ and $m = M$:

$$\alpha_{s1}^{(\mathrm{e})} = -\frac{1}{D^{(\mathrm{e})}}\left[r\omega^2-\frac{1}{r^2}M(1+n)\right],\tag{5.24}$$

$$\alpha_{s0}^{(\mathrm{e})} = -\frac{1}{D^{(\mathrm{e})}}\left[(n+1+\frac{1}{r}M)\omega^2-\frac{1}{r^4}M(1+n)(r-2M)\right],\tag{5.25}$$

with $D^{(\mathrm{e})}$ being:

$$D^{(\mathrm{e})} = \omega^4+\frac{1}{r^4}M(-7M+4r)\omega^2-\frac{2}{r^7}M^2(1+n)(r-2M).\tag{5.26}$$

VI. EQUATIONS OF MOTION IN H-S APPROACH

With the expressions of F and dF/dr_* derived above, we manage to rewrite the AAKS formalism in terms of the fields H and S, which is useful in decoupling matter and spacetime in oscillations of compact stars.

A. Equations inside the Star

The system inside the star can be described by two coupled second-order ODEs,

$$\frac{d^2S}{dr_*^2} - V^S_{f0}\alpha_{s1}\frac{dS}{dr_*} + (\omega^2 - V^S_{s0} - V^S_{f0}\alpha_{s0})S = V^S_{f0}(\alpha_{h1}\frac{dH}{dr_*} + \alpha_{h0}H), \tag{6.1}$$

$$\frac{d^2H}{dr_*^2} + (V^H_{h1} - V^H_{f1}\beta_{h1} - V^H_{f0}\alpha_{h1})\frac{dH}{dr_*} + (\frac{\omega^2}{C_s^2} - V^H_{h0} - V^H_{f1}\beta_{h0} - V^H_{f0}\alpha_{h0})H$$
$$= (V^H_{s1} + V^H_{f1}\beta_{s1} + V^H_{f0}\alpha_{s1})\frac{dS}{dr_*} + (V^H_{s0} + V^H_{f1}\beta_{s0} + V^H_{f0}\alpha_{s0})S, \tag{6.2}$$

where the α and β coefficients are defined in (5.14)-(5.21), and

$$V^S_{s0} = \frac{2e^\nu}{r^3}\left[(n+1)r - 2\pi r^3(\rho+3p) - m\right], \tag{6.3}$$

$$V^S_{f0} = -\frac{4e^{2\nu}}{r^5}\left[\frac{(m+4\pi pr^3)^2}{r-2m} + 4\pi\rho r^3 - 3m\right], \tag{6.4}$$

$$V^H_{h1} = \frac{e^{(\nu+\lambda)/2}}{r^2}\left[(m+4\pi pr^3)(1-\frac{1}{C_s^2}) + 2(r-2m)\right], \tag{6.5}$$

$$V^H_{h0} = \frac{2e^\nu}{r^2}\left[n+1-2\pi r^2(\rho+p)(3+\frac{1}{C_s^2})\right], \tag{6.6}$$

$$V^H_{f1} = (m+4\pi pr^3)(1-\frac{1}{C_s^2})\frac{e^{(\lambda+\nu)/2}}{2r^3}, \tag{6.7}$$

$$V^H_{f0} = \frac{e^\nu}{r^2}\left[\frac{2(m+4\pi pr^3)^2}{r^2(r-2m)}\frac{1}{C_s^2} - \frac{m+4\pi pr^3}{2r^2}(1-\frac{1}{C_s^2}) - 4\pi r(3p+\rho)\right], \tag{6.8}$$

$$V^H_{s1} = -(m+4\pi pr^3)(1-\frac{1}{C_s^2})\frac{e^{(\lambda-\nu)/2}}{2r}, \tag{6.9}$$

$$V^H_{s0} = \frac{(m+4\pi pr^3)^2}{r^2(r-2m)}(1+\frac{1}{C_s^2}) - \frac{m+4\pi pr^3}{2r^2}(1-\frac{1}{C_s^2}) - 4\pi r(3p+\rho). \tag{6.10}$$

The S-equation (6.1) can be recast into a much nicer form with the following transformation:

$$S(\omega, r_*) = q(\omega, r_*)\widetilde{S}(\omega, r_*), \tag{6.11}$$

where $q(\omega, r_*)$ is defined as:

$$q(\omega, r_*) = \exp\left[\frac{1}{2}\int_0^{r_*} V^S_{f0}(r'_*)\alpha_{s1}(\omega, r'_*)dr'_*\right]. \tag{6.12}$$

In terms of the transformed variable $\widetilde{S}(\omega, r_*)$, Eq. (6.1) becomes:

$$\frac{d^2\widetilde{S}}{dr_*^2} + [\omega^2 - \widetilde{V}^{(\mathrm{i})}(\omega, r_*)]\widetilde{S} = \widehat{O}_{\widetilde{S}}H, \tag{6.13}$$

with

$$\widetilde{V}^{(\mathrm{i})}(\omega, r_*) = V^S_{s0}(r_*) + V^S_{f0}(r_*)\alpha_{s0}(\omega, r_*) - \frac{1}{2}\frac{d}{dr_*}\left[V^S_{f0}(r_*)\alpha_{s1}(\omega, r_*)\right] + \frac{1}{4}\left[V^S_{f0}(r_*)\alpha_{s1}(\omega, r_*)\right]^2, \tag{6.14}$$

$$\widehat{O}_{\widetilde{S}} = \frac{1}{q(\omega, r_*)}V^S_{f0}(r_*)\left[\alpha_{h1}(\omega, r_*)\frac{d}{dr_*} + \alpha_{h0}(\omega, r_*)\right]. \tag{6.15}$$

The H-equation (6.2) can also be written in terms of $\widetilde{S}$,

$$\frac{d^2H}{dr_*^2} + (V^H_{h1} + \delta V^H_{h1})\frac{dH}{dr_*} + (\frac{\omega^2}{C_s^2} - V^H_{h0} - \delta V^H_{h0})H = \widehat{O}_H\widetilde{S}, \tag{6.16}$$

with

$$\begin{aligned}
\delta V^H_{h1} &= -V^H_{f1}\beta_{h1} - V^H_{f0}\alpha_{h1}, && (6.17)\\
\delta V^H_{h0} &= V^H_{f1}\beta_{h0} + V^H_{f0}\alpha_{h0}, && (6.18)\\
\widehat{O}_H &= (V^H_{s1} + V^H_{f1}\beta_{s1} + V^H_{f0}\alpha_{s1})\left[\frac{dq(\omega,r_*)}{dr_*} + q(\omega,r_*)\frac{d}{dr_*}\right] + (V^H_{s0} + V^H_{f1}\beta_{s0} + V^H_{f0}\alpha_{s0})\, q(\omega,r_*). && (6.19)
\end{aligned}$$

B. Equations outside the Star

Since the functions $\alpha_{h1}(\omega,r_*)$ and $\alpha_{h0}(\omega,r_*)$ both contain a $(\rho+p)$ factor, they vanish outside the star. Therefore, the system reduces into one second-order ODE in S outside the star:

$$\frac{d^2S}{dr_*^2} - V^{S(\mathrm{e})}_{f0}\alpha^{(\mathrm{e})}_{s1}\frac{dS}{dr_*} + (\omega^2 - V^{S(\mathrm{e})}_{s0} - V^{S(\mathrm{e})}_{f0}\alpha^{(\mathrm{e})}_{s0})S = 0, \tag{6.20}$$

where

$$\begin{aligned}
V^{S(\mathrm{e})}_{s0} &= \frac{2}{r^4}(r-2M)[(1+n)r - M], && (6.21)\\
V^{S(\mathrm{e})}_{f0} &= \frac{4}{r^7}(r-2M)M(3r-7M), && (6.22)
\end{aligned}$$

and the $\alpha^{(\mathrm{e})}$ coefficients are defined in (5.24) and (5.25).

As a counterpart of (6.13), $\widetilde{S}$ satisfies

$$\frac{d^2\widetilde{S}}{dr_*^2} + [\omega^2 - \widetilde{V}^{(\mathrm{e})}(\omega,r_*)]\widetilde{S} = 0, \tag{6.23}$$

where

$$\begin{aligned}
\widetilde{V}^{(\mathrm{e})}(\omega,r_*) &= V^{S(\mathrm{e})}_{s0}(r_*) + V^{S(\mathrm{e})}_{f0}(r_*)\alpha^{(\mathrm{e})}_{s0}(\omega,r_*) - \frac{1}{2}\frac{d}{dr_*}\left[V^{S(\mathrm{e})}_{f0}(r_*)\alpha^{(\mathrm{e})}_{s1}(\omega,r_*)\right] + \frac{1}{4}\left[V^{S(\mathrm{e})}_{f0}(r_*)\alpha^{(\mathrm{e})}_{s1}(\omega,r_*)\right]^2,\\
&= \frac{2(r-2M)}{[D^{(\mathrm{e})}r^9]^2}\Big\{2M^4(n+1)^2(2M-r)[3M^2 - 2r^2(n+1) + 2Mr(2n+1)]\\
&\quad -M^3r^3(n+1)\omega^2[126M^3 - 246M^2r - Mr^2(8n-131) + 4r^3(n-5)]\\
&\quad -M^2r^6\omega^4[147M^3 - M^2r(97n+258) - 4Mr^2(2n^2-17n-33) + r^3(4n^2-11n-21)]\\
&\quad -Mr^{10}\omega^6[56M^2 - 51Mr - 2r^2(n-5)] + r^{14}\omega^8[(n+1)r - M]\Big\} && (6.24)
\end{aligned}$$

and $\widetilde{S}$ is related to S through

$$S(\omega,r_*) = q^{(\mathrm{e})}(\omega,r_*)\widetilde{S}(\omega,r_*), \tag{6.25}$$

with $q^{(\mathrm{e})}(\omega,r_*)$ given by:

$$q^{(\mathrm{e})}(\omega,r_*) = q(\omega,R_*)\exp\left[\frac{1}{2}\int_{R_*}^{r_*} V^{S(\mathrm{e})}_{f0}(r'_*)\alpha^{(\mathrm{e})}_{s1}(\omega,r'_*)dr'_*\right]. \tag{6.26}$$

Hereafter R_* is the tortoise coordinate of the stellar surface $r = R$. From the forms of $V^{S(\mathrm{e})}_{f0}$ and $\alpha^{(\mathrm{e})}_{s1}$, it is easy to see that $q^{(\mathrm{e})}$ approaches a constant at the spatial infinity.

It is interesting to note that the Zerilli function is expressible solely in terms of S (or equivalently $\widetilde{S}$) in the H-S approach as H is identically zero outside the star, which reads as:

$$\begin{pmatrix} Z \\ dZ/dr_* \end{pmatrix} = \begin{pmatrix} Z_{11}(\omega,r) & Z_{12}(\omega,r) \\ Z_{21}(\omega,r) & Z_{22}(\omega,r) \end{pmatrix} \begin{pmatrix} S \\ dS/dr_* \end{pmatrix}, \tag{6.27}$$

with

$$Z_{11} = \frac{1}{D^{(\mathrm{e})}}\left\{\frac{M(r-2M)\,[9M+(-3+n)\,r]}{(3M+nr)r^4}+\omega^2\right\}, \tag{6.28}$$

$$Z_{12} = \frac{1}{D^{(\mathrm{e})}}\frac{M[3M-(3+n)r]}{(3M+nr)r^2}, \tag{6.29}$$

$$Z_{21} = \frac{M\left[-9M^2+9Mr+n(n+3)r^2\right]\omega^2}{D^{(\mathrm{e})}r^2(3M+nr)^2} - \frac{M(r-2M)}{D^{(\mathrm{e})}r^6(3M+nr)^2}\left[54M^3+3(14n-3)M^2r+6n(3n+1)Mr^2+n(n+1)(2n-3)r^3\right], \tag{6.30}$$

$$Z_{22} = \frac{1}{D^{(\mathrm{e})}}\left[\frac{3M(r-2M)\left[3M^2+2nMr+n(n+1)r^2\right]}{r^4(3M+nr)^2}+\omega^2\right]. \tag{6.31}$$

Therefore, Eq. (6.20) or Eq. (6.23) can be considered as an appropriate replacement for the Zerilli equation in the current theory. Likewise, S or $\widetilde{S}$ can in fact play the role of the Zerilli function Z. For example, QNMs can be defined by the imposing the outgoing wave boundary condition on S (or $\widetilde{S}$).

To sum up, in the H-S approach of the AAKS formalism a pulsating NS is governed by two coupled second-order equations, (6.13) and (6.16), inside the star, and one single second-order equation (6.23) outside. The polar QNMs can be located by imposing the regularity boundary condition at the star center and the outgoing boundary condition at the spatial infinity. In the following discussion, we will show that the H-S approach achieved here forms the basis of a decoupling scheme readily leading to accurate CA and ICA for p-modes and w-modes, respectively.

VII. COWLING APPROXIMATION

The p-mode pulsation is one important type of fluid modes, whose restoring force comes mainly from pressure, and has been well studied for non-relativistic stars under the CA [29]. Here, based on the H-S approach developed above, we propose a relativistic version of CA for p-modes of compact stars. We will also see that such a CA scheme could also provide qualitatively correct description for the f-mode

A. Fluid equation

It is well known that the eigenfrequencies of the p-modes of a compact star are characterized by dramatically small imaginary parts. The smallness of the imaginary part implies the tininess of GWs emitted in p-modes. Therefore, the influence of the variable S on the fluid motion is expected to be negligible when calculating p-mode frequencies. With this in mind, we propose that the approximate eigenfrequencies of p-modes can be calculated with the following equation:

$$\frac{d^2H}{dr_*^2}+(V_{h1}^H+\delta V_{h1}^H)\frac{dH}{dr_*}+(\frac{\omega^2}{C_s^2}-V_{h0}^H-\delta V_{h0}^H)H=0, \tag{7.1}$$

which is obtained by neglecting the $\widetilde{S}$ part in (6.16). Furthermore, we find that for p-mode oscillations the effects of δV_{h1}^H and δV_{h0}^H are pretty small. Then, Eq. (7.1) can be further simplified into:

$$\frac{d^2H}{dr_*^2}+V_{h1}^H\frac{dH}{dr_*}+(\frac{\omega^2}{C_s^2}-V_{h0}^H)H=0. \tag{7.2}$$

The above equation can also be derived by simply neglecting the S, dS/dr_*, F and dF/dr_* terms in (4.9). Unlike δV_{h0}^H and δV_{h1}^H, which are frequency-dependent, V_{h0}^H and V_{h1}^H are both frequency-independent. Notwithstanding the fact that Eq. (7.1) properly accounts for the effects of F on the fluid motion and is proved to be the best approximation for p-modes in the following discussion, the mathematical structure of (7.2) is more desirable and leads to a complete orthonormal set of eigenfunctions.

B. Boundary conditions

Under CA (7.2), p-mode oscillations are determined from (7.2) by imposing proper boundary conditions on H at $r=0$ and $r=R$. Following directly from (4.22), $H \approx h_0 r_*^l$ near the star center. Assuming that the EOS acquires the polytropic form $p \propto \rho^{1+1/N}$ with the polytropic index $N>1$ near the star surface, we show that (see Appendix A)

$$C_s^2 \sim A_c(R_* - r_*), \tag{7.3}$$

with $A_c = M/(R^2 N)$. Consequently, the H-equation near the stellar surface reduces asymptotically to

$$\frac{d^2H}{dr_*^2} - \frac{M}{R^2 A_c(R_* - r_*)}\frac{dH}{dr_*} + \frac{\omega^2}{A_c(R_* - r_*)}H = 0. \tag{7.4}$$

There are two independent solutions for this equation. The first one is bounded and behaves as:

$$H \propto \left[1 + \omega^2 R^2 (r_* - R_*)/M\right]. \tag{7.5}$$

The other one is proportional to $(R_* - r_*)^{1-N}$ and is unbounded (also see Appendix A). As argued previously, H is finite at the stellar surface and therefore follows (7.5).

The single H-equation (7.2) together with two boundary conditions (4.22) and (7.5) form an eigenvalue problem. The approximate eigenfrequencies of polar p-modes under the CA can then be located.

C. Completeness and orthogonality of p-modes

The eigenvalue problem of the purely matter-based H-equation (7.2) is actually a Sturm-Liouville eigenvalue problem, as it can be cast into the following standard form:

$$\frac{d}{dr_*}\left[P(r_*)\frac{dH}{dr_*}\right] - Q(r_*)H = -\omega^2 \Lambda(r_*) H, \tag{7.6}$$

with

$$\begin{aligned} P(r_*) &= e^{\nu}(\rho+p)r^2, && (7.7)\\ Q(r_*) &= 2e^{2\nu}(\rho+p)\left[n+1-2\pi r^2(\rho+p)(3+1/C_s^2)\right], && (7.8)\\ \Lambda(r_*) &= e^{\nu}(\rho+p)r^2/C_s^2. && (7.9)\end{aligned}$$

There are a series of real eigenfrequencies $\{\sigma_n\}$ and the corresponding normalized eigenfunctions $\{H_n(r_*)\}$, which are defined by the boundary conditions derived above, namely (i) H is regular at $r_*=0$, (ii) $(dH/dr_*)/H = R^2\omega^2/M$ at $r=R_*$. Following straightforwardly from the standard theory of Sturm-Liouville system and the fact that $\Lambda(r=0) = 0 = \Lambda(r=R)$, the normalized eigenfunctions $\{H_n(r_*)\}$ form a complete orthogonal set obeying the completeness and orthogonality relationships (see, e.g., [33]):

$$\begin{aligned} \sum_n \Lambda(r_*)H_n(r_*)H_m(r'_*) &= \delta(r_* - r'_*); && (7.10)\\ \int_0^{R_*} dr_* \Lambda(r_*)H_n(r_*)H_m(r_*) &= \delta_{nm}. && (7.11)\end{aligned}$$

D. Numerical results and discussions

To gauge the accuracy of the CA developed (or ICA as discussed later) in the present paper, we use EOS A [42] to construct a NS with compactness $\mathcal{C} = 0.27$ and central density $\rho_c = 2.227 \times 10^{-3}$ and evaluate the QNMs of the star. Table I shows the frequencies of the leading fluid modes obtained from exact calculation and CA (7.2) developed above. We see that the lowest eigenfrequency σ_0 of (7.6) provides us an approximate value of the real part of f-mode frequency, $\omega_{\mathrm{r},f}$, with a percentage error of around 20%. For $n \geq 1$, the eigenfrequency σ_n yields an accurate approximation for the real part of the frequency of the n-th p-mode, $\omega_{\mathrm{r},n}$. The percentage error is usually much less than 1% and decreases with increasing n (see Table I).

The comparison between H-function obtained from (7.2) and the exact H-function is shown in Fig. 1. Both of them are normalized so that

$$\int_0^{R_*} dr_* \Lambda(r_*) |H|^2 = 1\,. \tag{7.12}$$

Besides, we assume that the exact H-function is real at $r = R$. The numerical results again confirm that the CA proposed here is very accurate for p-modes. However, the discrepancy between the approximate and the exact wave functions grows larger for f-mode, which implies that the coupling between spacetime (i.e., F and S) and matter is negligible for p-modes, but is not so weak for f-mode. Actually, from the form of (6.2), it is clearly seen that the accuracy of the CA (7.2), would decrease if $D(r, \omega)$ is small. As shown in Fig. 2, where D is plotted against r/R for different values of ω, D decreases with decreasing ω. In fact, for $\omega = \sigma_0$ (i.e., the f-mode frequency under CA), D vanishes at $r/R = 0.947$. Therefore, the omission of the influence of spacetime (i.e., F and S) on oscillations of matter field may lead to appreciable errors at low frequencies. On the other hand, we observe that there are exactly n nodes (excluding the origin) in the wave function H_n, in agreement with standard theory of the Sturm-Liouville eigen-system (see, e.g., [43]).

We also note that the high accuracy of the CA (7.2) is generic and independent of the EOS and the compactness of the NS. Tables II and III show the exact complex eigenfrequencies $(\omega_{\rm r}, \omega_{\rm i})$ and approximate real eigenfrequencies σ_n obtained from CA scheme (7.2) for the leading fluid modes of NSs with Sly EOS [44, 45] and polytropic EOS: $p = 100\rho^{1.8}$, respectively. It is clearly seen that (7.2) indeed leads to accurate numerical results for the p-modes irrespective of the EOS and the compactness of the star. Again the accuracy improves with increasing frequency, reminding us of the insignificance of the coupling between spacetime and matter in high order p-modes. As for the f-modes, we note that the performance of CA (7.2) improves for polytropic stars with low compactness.

For the purpose of comparison, approximate eigenfrequencies σ_n obtained from (7.1) are also listed in Tables I, II and III. While Eq. (7.1) in general can yield almost exact numerical results for p-modes because it has properly taken the influence of F on H into consideration, it completely fails to locate approximate position of the f-mode. The inability of (7.1) to handle the f-mode can be understood as a direct consequence of the fact that D could vanish at certain positions inside the star at frequencies close to the f-mode frequency (see Fig. 2).

VIII. INVERSE-COWLING APPROXIMATION FOR w-MODE

A. Spacetime oscillations

From the nature of polar w-mode oscillations, we know that the coupling between fluid motion and spacetime oscillations is very weak (see, e.g., [32]). Therefore, fluid motion is hardly involved in w-mode and accordingly the contribution of the H and dH/dr_* terms in (6.1) is negligible. Hence, the polar w-mode system can be described by a single second-order equation:

$$\frac{d^2 S}{dr_*^2} - V_{f0}^S \alpha_{s1} \frac{dS}{dr_*} + (\omega^2 - V_{s0}^S - V_{f0}^S \alpha_{s0}) S = 0, \tag{8.1}$$

where V_{s0}^S, V_{f0}^S, α_{s1} and α_{s0} take the forms (6.3), (6.4), (5.14) and (5.15) inside the star, and take the forms (6.21), (6.22), (5.24) and (5.25) outside the star. As the fluid variable H is eliminated, the method is another form of ICA [31, 32]. To our knowledge, this is the first time that a single second-order equation for polar w-modes has been proposed and verified numerically (see the following discussion).

The eigenvalue system can be simplified a lot after turning into the $\widetilde{S}$ representation. The spacetime oscillations are described by the following equation in $\widetilde{S}$ representation,

$$\widehat{L}_{\widetilde{S}}\, \widetilde{S}(r_*) = \left[\frac{d^2}{dr_*^2} + \omega^2 - \widetilde{V}(\omega, r_*) \right] \widetilde{S}(r_*) = 0, \tag{8.2}$$

where $\widetilde{V}$ takes the form $\widetilde{V}^{(\rm i)}$ (6.14) and $\widetilde{V}^{(\rm e)}$ (6.24) inside and outside the star, respectively.

B. Numerical results and discussions

The spacetime w-modes can be located by solving (8.1) or equivalently (8.2) with the outgoing boundary condition. The numerical results of ICA proposed here (the columns labelled with ICA) are compared with the exact polar

w-mode eigenfrequencies in Table IV, where the exact and approximate QNM frequencies of the leading twelve polar w-modes (including two $w_{\rm II}$ modes) for the above-mentioned EOS A NS ($\mathcal{C}=0.27$) are listed. The agreement between the exact frequencies and the ICA frequencies is almost perfect, especially for $\omega_{\rm r}$ and higher order modes.

C. High-frequency ICA

We note that both α_{s0} and α_{s1} go as $1/\omega^2$ at high frequencies. Thus, we expect that w-mode QNMs with high frequencies can be approximately located by omitting terms proportional to α_{s0} and α_{s1} in (8.1), namely,

$$\omega^2 S + \frac{d^2 S}{dr_*^2} + \frac{2e^\nu}{r^3}\left[2\pi r^3(\rho+3p) + m - (n+1)r\right] S = 0. \tag{8.3}$$

Such a scheme is termed as the high-frequency inverse-Cowling approximation (HFICA) in the present paper. As shown in Table IV, HFICA works well in the high frequency regime. In fact, HFICA can successfully explain the high-frequency asymptotic behavior of polar w-modes of compact stars and an iterative method has be formulated to improve the accuracy of HFICA for the low-order modes [46].

IX. CONCLUSION AND DISCUSSION

In this paper, based on the AAKS formalism for pulsations of compact stars [23], we establish an equivalent description consisting of two coupled second-order equations in two variables, namely H and S, measuring relativistic dynamics of matter and spacetime, respectively. In addition to simplification of the theory proposed in [23], the H-S approach developed here readily leads to CA (ICA) with unprecedented accuracies for polar p-modes (w-modes) when the two second-order equations are decoupled. Besides, under the CA the wave functions H_n ($n=0,1,2,\ldots$) of the fluid modes (including the f-mode and the p-modes) are shown to form a complete orthonormal set, which paves the way for further developing perturbation schemes to improve the accuracies of the CA and to include the effect of GW radiation damping.

We note that there are other theories for pulsations of compact stars which also result in two coupled second-order equations. For example, in the LMI formalism [21] the metric variable $\bar{H}_0(r,t)$ and the fluid variable $\delta U(r,t) \equiv \delta p(r,t)/(\rho+p)+\bar{H}_0(r,t)/2$ are used to formulate a relativistic theory for stellar pulsations. Similar to the development of the present paper, the two coupled second-order equations in the LMI formalism can be decoupled by adopting the approximations $\bar{H}_0(r,t)=0$ and $\delta U=0$, leading to another version of CA and ICA, respectively. The CA developed in the LMI formalism was applied to find the frequency of the f-mode of a polytropic NS with $M=1.4M_\odot$ and the percentage error was about 20% for the case $l=2$ [21], which is similar to ours. We also compute the QNM frequencies of the leading fluid modes of the EOS A star considered above ($\mathcal{C}=0.27$) with the CA of the LMI formalism and list them in Table I. It is seen that the CA in the LMI formalism can yield accurate frequencies for the p-modes. On the other hand, the frequencies of the leading polar w-modes obtained from the ICA scheme suggested in the LMI formalism (the columns labelled with Ref. [21]) are listed in the Table IV. It is clear, from the numerical results, that the performance of the ICA scheme mentioned in [21] is far from satisfactory.

The success of the CA and ICA established in the present paper based on the AAKS formalism [23] is attributable to the selection of independent variables, namely H and S. While the former is proportional to the change in the Eulerian pressure, i.e., a proper Newtonian quantity, the latter vanishes in the weak field limit. Thus, the interplay between H and S in fluid modes and spacetime modes is expected to be negligible, which is numerically verified in the above discussion. According to our experience, the choice of the independent variables H and S is the most crucial factor leading to the high precision of the ICA scheme developed here. We note that several other schemes for CA (see, e.g., [30] and references therein) and ICA [31] based on the LD formalism have been proposed. Yet, the accuracy of these schemes is worse than that achieved in the present paper.

Lastly, as F is proportional to the change in the Newtonian potential in the weak field limit, it can also reveal the motion of matter indirectly and replace the role of H. In fact, one can also formulate another equivalent description for stellar pulsation using the variables F and S to develop the F-S approach based on the AAKS formalism (see Appendix B for detailed discussion). Similar to the case in the LMI formalism mentioned above [21], such an F-S approach can yield valid CA with accuracies worse than those obtained from (7.2) and (7.1), but fails to predict accurate polar w-modes. The ICA in the F-S approach is identical to the HFICA in the H-S approach and hence works well only in the high frequency limit (see Appendix B and Table IV). Instead, according to our experience, Eq. (8.1), or equivalently (8.2), can locate all polar w-modes including the low-frequency $w_{\rm II}$ modes with excellent numerical accuracies. In a nutshell, the H-S approach proposed here based on the AAKS formalism [23] successfully leads to feasible and accurate CA and ICA schemes in a unified framework.

Acknowledgments

This work is supported in part by the Hong Kong Research Grants Council (Grant No: 401807) and the direct grant (Project ID: 2060330) from the Chinese University of Hong Kong. We thank L.M. Lin, H.K. Lau, P.O. Chan and T.K. Chan for helpful discussions.

Appendix A: Boundary conditions at the stellar surface

Near the stellar surface, we can assume that the behavior of the matter component is well captured by the polytropic EOS $p = \kappa\rho^{\gamma}$, where $\kappa > 0$ is a constant and $\gamma \equiv 1 + 1/N$, with N being the effective polytropic index there. Without loss of generality and also due to the behavior of nuclear matter at low densities, we assume that $N > 1$. It follows directly from the TOV equations that [17]

$$p = \alpha\zeta^{N+1} + \alpha_1\zeta^{N+2} + \cdots, \tag{A1}$$

$$\rho = \beta\zeta^{N} + \beta_1\zeta^{N+1} + \cdots, \tag{A2}$$

with $\zeta \equiv R - r$ and

$$\alpha = \kappa\left[\frac{M}{\kappa R(R-2M)(N+1)}\right]^{N+1}, \tag{A3}$$

$$\alpha_1 = \frac{[M - 2(N+1)(M-R)]M^{N+1}}{2\kappa^N[R(R-2M)]^{N+2}(N+1)^{N+1}}, \tag{A4}$$

$$\beta = \left[\frac{M}{\kappa R(R-2M)(N+1)}\right]^{N}, \tag{A5}$$

$$\beta_1 = \frac{N[M - 2(N+1)(M-R)]M^{N}}{2\kappa^N[R(R-2M)(N+1)]^{N+1}}. \tag{A6}$$

Similarly, it can also be shown that

$$m(r) = \int_0^r 4\pi r'^2\rho(r')dr' = M - 4\pi\beta R^2\zeta^{N+1} + \cdots, \tag{A7}$$

$$e^{\nu} = \left(1 - \frac{2M}{R}\right)\left[1 - \frac{2M\zeta}{R(R-2M)}\right] + \cdots, \tag{A8}$$

$$e^{\lambda} = \frac{R}{R-2M}\left[1 + \frac{2M\zeta}{R(R-2M)}\right] + \cdots, \tag{A9}$$

$$C_s^2 = \frac{M}{R(R-2M)N}\zeta + \frac{M[M - 2(N+1)(M-R)]}{2N(N+1)R^2(R-2M)^2}\zeta^2 + \cdots, \tag{A10}$$

$$\frac{1}{C_s^2} = \frac{NR^2}{M\zeta_*} + 3N - \frac{NR}{M} - \frac{N}{2(N+1)} + \cdots, \tag{A11}$$

where

$$\begin{aligned}\zeta_* &\equiv \int_r^R \exp\left\{\left[\lambda(r') - \nu(r')\right]/2\right\} dr' \\ &= \frac{R\zeta}{R-2M}\left[1 + \frac{2M\zeta}{R(R-2M)}\right] + \cdots.\end{aligned} \tag{A12}$$

Now we consider the behavior of the wave functions H, S and F near the stellar surface. Since the coefficients in evolution equations (e.g., ρ, p and m) are expanded in powers of ζ and ζ^N, the wave functions near the surface are accordingly given by:

$$H = \zeta_*^{a}\zeta_*^{a'N}\sum_{m,n=0} h_{mn}\zeta_*^{m}\zeta_*^{nN}, \tag{A13}$$

$$S = \zeta_*^{b}\zeta_*^{b'N}\sum_{m,n=0} s_{mn}\zeta_*^{m}\zeta_*^{nN}, \tag{A14}$$

$$F = \zeta_*^{c}\zeta_*^{c'N}\sum_{m,n=0} f_{mn}\zeta_*^{m}\zeta_*^{nN}, \tag{A15}$$

where a, a', b, b', c, c', h_{mn}, s_{mn} and f_{mn} are constants to be determined from the AAKS equations and the boundary conditions at the stellar surface, and h_{00}, s_{00} and f_{00} are non-zero by definition. Based on the boundedness of $H(r = R)$ and the AAKS equations, we can show that all the powers a, a', b, b', c, and c' vanish.

In principle all the coefficients h_{mn}, s_{mn} and f_{mn} can be determined from the modified Hamiltonian constraint (4.14) and the AAKS equations (4.7) and (4.8). However, for the sake of the present paper, we only need to find the relevant expansions to leading few orders as shown below:

$$S \approx s_{00} + s_{10}\zeta_* + s_{20}\zeta_*^2 + s_{01}\zeta_*^N + s_{11}\zeta_*^{N+1}, \tag{A16}$$

$$F \approx f_{00} + f_{10}\zeta_* + f_{20}\zeta_*^2 + f_{01}\zeta_*^N + f_{11}\zeta_*^{N+1}, \tag{A17}$$

$$H \approx h_{00} + h_{10}\zeta_*, \tag{A18}$$

with

$$s_{10} = -\frac{R^7\omega^4 + 4MR^4\omega^2 - M^2R(2+2n+7R^2\omega^2) + 4M^3(n+1)}{R^5[R^3\omega^2 - M(n+1)]} f_{00} \tag{A19}$$

$$-\frac{(n+1)R^4\omega^2 + MR(R^2\omega^2 - 1 - n) + 2M^2(n+1)}{R^2[R^3\omega^2 - M(n+1)]} s_{00}, \tag{A20}$$

$$s_{01} = 0, \tag{A21}$$

$$s_{11} = 0, \tag{A22}$$

$$s_{20} = -\frac{1}{2}\left\{\omega^2 + \frac{2(R-2M)}{R^4}\left[M - (n+1)R\right]\right\} s_{00} - \frac{2M(R-2M)(7M-3R)}{R^7} f_{00}, \tag{A23}$$

$$f_{10} = \frac{M(n+1)(2M-R) + R^3[M - (n+2)R]\omega^2}{R^2[R^3\omega^2 - M(n+1)]} f_{00}$$

$$-\frac{(n+1)R(2M+nR) + R^4\omega^2}{R^3\omega^2 - M(n+1)} s_{00}, \tag{A24}$$

$$f_{01} = 0, \tag{A25}$$

$$f_{11} = -8\pi\beta\frac{(R-2M)^{N+1}}{R^{N-2}M(N+1)} h_{00}, \tag{A26}$$

$$f_{20} = -\frac{1}{2}\left\{\omega^2 + \frac{2(R-2M)}{R^4}\left[M - (n+1)R\right]\right\} f_{00} + \frac{(R-2M)}{R} s_{00}, \tag{A27}$$

$$\begin{aligned} h_{10} &= \frac{R}{2(R-2M)} s_{00} - \frac{R^2}{2(R-2M)} s_{10} + \frac{2M+R}{2R^3} f_{00} + \frac{1}{2R} f_{10} - \frac{R^2\omega^2}{M} h_{00} \\ &= \frac{2M(n+1)(2M-R)(M+R) + R^3[-13M^2 + (2n+9)MR - (n+1)R^2]\omega^2 + R^7\omega^4}{2(R-2M)R^3[M(n+1) - R^3\omega^2]} f_{00} \\ &\quad + \frac{(n+1)[6M^2 + 2(n-2)MR - nR^2] + R^3[3M + (n+1)R]\omega^2}{2(R-2M)[M(n+1) - R^3\omega^2]} s_{00} - \frac{R^2\omega^2}{M} h_{00}. \end{aligned} \tag{A28}$$

While the expressions for s_{01} (f_{01}), s_{11} (f_{11}), and s_{20} (f_{20}) are obtained by comparing the coefficients of ζ_*^{N-2}, ζ_*^{N-1} and ζ_*^0 terms, respectively, in Eq. (4.7) (Eq. (4.8)), those for s_{10}, f_{10}, h_{10} can be found from the coefficients of ζ_*^0, ζ_*^1 and ζ_*^{N+1} terms in the modified Hamiltonian constraint (4.14). On the other hand, the values of s_{00}, f_{00}, h_{00} can be considered as the three degrees of freedom of the AAKS equations and other expansion coefficients are all expressible in terms of them. Together with the two degrees of freedom arising at the origin, there are five degrees of freedom for the solution of the AAKS equations.

With the expansions (A16) and (A17) obtained above, we can match them with the ones obtained from numerically integrating (4.7) and (4.8) outward from the origin at a point near the surface of the star where the expansions are valid. The continuity of S, dS/dr_*, F and dF/dr_* lead to four independent equations, thus reducing the degree of freedom of the solution from five to one. As a result, the solutions for S, F and H (evaluated from the modified Hamiltonian constraint) are determined up to a common multiplicative constant.

On the other hand, there is another group of solution with divergent fluid variable H at the stellar surface:

$$S = \sum_{m,n=0} \tilde{s}_{mn}\zeta_*^m \zeta_*^{nN}, \tag{A29}$$

$$F = \sum_{m,n=0} \tilde{f}_{mn}\zeta_*^m \zeta_*^{nN}, \tag{A30}$$

$$H = \zeta_*^{-N+1} \sum_{m,n=0} \tilde{h}_{mn}\zeta_*^m \zeta_*^{nN}. \tag{A31}$$

To the leading few orders, the wave functions are given by:

$$S = \tilde{s}_{00} + \tilde{s}_{10}\zeta_* + \tilde{s}_{20}\zeta_*^2, \tag{A32}$$

$$F = \tilde{f}_{00} + \tilde{f}_{10}\zeta_* + \tilde{f}_{20}\zeta_*^2, \tag{A33}$$

$$H = \tilde{h}_{00}\zeta_*^{-N+1}, \tag{A34}$$

with

$$\tilde{s}_{10} = s_{10}, \tag{A35}$$

$$\tilde{s}_{01} = s_{01} = 0, \tag{A36}$$

$$\tilde{s}_{20} = s_{20}, \tag{A37}$$

$$\tilde{s}_{11} = s_{11} = 0, \tag{A38}$$

$$\tilde{f}_{10} = f_{10}, \tag{A39}$$

$$\tilde{f}_{01} = f_{01} = 0, \tag{A40}$$

$$\tilde{f}_{20} = f_{20} - 4\pi\beta\frac{(R-2M)^{N+1}}{R^{N-2}M}\tilde{h}_{00}. \tag{A41}$$

Appendix B: F-S Approach

1. Formalism

As both F and H could reveal the Newtonian aspect of stellar pulsations, it is natural to expect that an alternative approach using F and S to study stellar pulsations is possible. In fact, it is straightforward to make use of the time-independent Hamiltonian constraint (4.13) to eliminate the matter field $H(r_*)$ in (4.8) [39]:

$$\begin{aligned}&\frac{\omega^2 F}{C_s^2} + \frac{d^2F}{dr_*^2} - \left(1-\frac{1}{C_s^2}\right)\frac{e^{(\nu+\lambda)/2}}{r^2}(m+4\pi p r^3)\frac{dF}{dr_*} + \frac{e^\nu}{r^3}\left[4\pi r^3\left(3\rho+\frac{p}{C_s^2}\right) - m\left(1-\frac{3}{C_s^2}\right) - 2(n+1)r\right]F \\ &= \left(1-\frac{1}{C_s^2}\right) r e^{-(\nu+\lambda)/2}\frac{\partial S}{\partial r_*} + \left[2e^{-\lambda} + \left(1-\frac{1}{C_s^2}\right)(n+1) - 8\pi(p+\rho)r^2\right]S.\end{aligned} \tag{B1}$$

(4.7) and (B1) together form two coupled second-order equations in F and S, which can be used to locate QNMs of compact stars and lead to the so-called *F-S* approach in the present paper.

Similar to the *H-S* approach, the S terms in (B1) are expected to be negligible for fluid mode pulsations, yielding

$$\begin{aligned}&\frac{\omega^2 F}{C_s^2} + \frac{d^2F}{dr_*^2} - \left(1-\frac{1}{C_s^2}\right)\frac{e^{(\nu+\lambda)/2}}{r^2}(m+4\pi p r^3)\frac{dF}{dr_*} \\ &+\frac{e^\nu}{r^3}\left[4\pi r^3\left(3\rho+\frac{p}{C_s^2}\right) - m\left(1-\frac{3}{C_s^2}\right) - 2(n+1)r\right]F = 0.\end{aligned} \tag{B2}$$

When subjected to suitable boundary condition (as discussed in the following), (B2) could give good approximation to p-mode pulsations of realistic compact stars, thereby resulting in the CA for p-modes in the *F-S* approach.

On the other hand, we could also obtain the ICA for polar w-modes within the *F-S* approach by neglecting the F term in (4.7), i.e.,

$$\omega^2 S + \frac{d^2S}{dr_*^2} + \frac{2e^\nu}{r^3}\left[2\pi r^3(\rho+3p) + m - (n+1)r\right]S = 0. \tag{B3}$$

Interestingly, such an ICA is merely the HFICA in the H-S approach, which works well only for w-modes with high frequencies (see Table IV for the accuracies and validity of HFICA in the H-S approach). It reflects the fact that the metric variable F also participates in the dynamics of stellar pulsations and its contribution to (4.7) is non-negligible for w-modes with low frequencies.

2. Cowling approximation

Within the F-S approach, p-mode pulsations of compact stars in the CA can be located by solving (B2) with properly imposed boundary conditions at the two endpoints $r = 0$ and $r = R$. As usual, the regularity boundary condition $F \propto r_*^l$ holds around the origin. On the other hand, by assuming the polytropic EOS mentioned in Appendix A near the stellar surface, Eq. (B2) becomes

$$\frac{d^2F}{d\zeta_*^2} - \left(\frac{N}{\zeta_*} + A + \cdots\right)\frac{dF}{d\zeta_*} + \left(\frac{B}{\zeta_*} + C + \cdots\right)F = 0, \tag{B4}$$

with

$$\begin{aligned} A &= -\frac{M}{R^2}\left[N + 1 - \frac{NR}{M} + \frac{N}{2(N+1)}\right], && \text{(B5)}\\ B &= \frac{\omega^2R^2N}{M} + \frac{3N(R-2M)}{R^2}, && \text{(B6)}\\ C &= \omega^2\left[3N - \frac{NR}{M} - \frac{N}{2(N+1)}\right] - \frac{R-2M}{R^4}\left[M(1-9N) + \frac{3NM}{2(N+1)} - 2(n+1)R - 6NR\right]. && \text{(B7)} \end{aligned}$$

Correspondingly, there are two possible solutions near the surface, namely $F_a \sim f_0 + f_1\zeta_* + f_2\zeta_*^2 + \cdots$ and $F_b \sim f_{11}\zeta_*^{N+1} + \cdots$, where

$$\begin{aligned} f_1 &= [\frac{\omega^2R^2}{M} + \frac{3(R-2M)}{R^2}]f_0; && \text{(B8)}\\ f_2 &= \frac{f_0}{N-1}\left\{\frac{2M-R}{R^4}[M(3N-1) - R(1+7N)] - \frac{\omega^2}{2M}[M(8N-1) - 4NR] + \frac{NR^4\omega^4}{2M^2}\right\}. && \text{(B9)} \end{aligned}$$

In the CA, S is neglected and hence the modified Hamiltonian constraint (4.14) leads to

$$8\pi re^{\nu}(\rho + p)H = \frac{e^{(\nu+\lambda)/2}}{r^2}(m + 4\pi r^3p)\frac{dF}{dr_*} + \omega^2F + \frac{e^{\nu}}{r^3}(4\pi r^3p + 3m)F. \tag{B10}$$

By inserting the asymptotic forms F_a and F_b into the above equation, the corresponding expansion of H at the stellar surface can be obtained, namely, $H_a \sim h_0\zeta_*^{-(N+1)}$ and $H_b \sim h_{00}$, where

$$\begin{aligned} &\left\{\frac{2(2M-R)}{R^4}[2NM - (n+2)R + N(n-6)R] + \frac{N\omega^2}{M}(4R-7M) + \frac{N\omega^4R^4}{M^2}\right\}f_0 \\ &= -8\pi\beta\frac{(R-2M)^{N+1}}{R^{N-1}}\frac{N(N-1)}{M}h_0, && \text{(B11)}\\ f_{11} &= -8\pi\beta\frac{(R-2M)^{N+1}}{R^{N-2}M(N+1)}h_{00}. && \text{(B12)} \end{aligned}$$

As the fluid variable H is bounded, the correct boundary condition near the stellar surface is $F \sim f_{11}\zeta_*^{N+1} + \cdots$.

Upon imposing the two boundary conditions mentioned above on the solution of (B2), we can find the approximate values of the real part of the eigenfrequencies of p-modes. However, as will be discussed and explained in the following, we cannot locate the f-mode oscillation under the CA in this approach.

3. Completeness and orthogonality of p-modes

Starting from CA (B2) in the F-S approach, we can get a standard Sturm-Liouville equation in terms of the variable $G \equiv F/p$,

$$\frac{d}{dr_*}\left[P_G(r_*)\frac{dG}{dr_*}\right] - Q_G(r_*)G + \omega^2\Lambda_G(r_*)G = 0, \tag{B13}$$

with

$$\begin{aligned} P_G(r_*) &= \frac{p^2}{e^{\nu}(\rho+p)}, && \text{(B14)} \\ Q_G(r_*) &= -\frac{p}{2r(r-2m)}\left[16\pi mr(\rho+3p)-64\pi^2r^4p^2-8\pi r^2(\rho+p)+1\right]+\frac{(r-2m)p}{2r^3} \\ &\quad -\frac{p^2}{r^3(\rho+p)}\left[4\pi r^3(3\rho+\frac{p}{C_s^2})-m(1-\frac{3}{C_s^2})-2(n+1)r\right], && \text{(B15)} \\ \Lambda_G(r_*) &= \frac{p^2}{e^{\nu}(\rho+p)C_s^2}. && \text{(B16)} \end{aligned}$$

We note that the coefficients $P_G(r_*)$ and $\Lambda_G(r_*)$ are strictly positive, as required for a normal Sturm-Liouville equation (see, e.g., [43]). Besides, Λ_G vanishes at both $r = 0$ and $r = R$. Following directly from the physical boundary condition on F, namely, $F \sim \zeta_*^{N+1}$ near the stellar surface, G tends to a finite constant there. In fact, by expanding (B13) about the regular singular point at $r_* = R_*$, we find the corresponding boundary condition for G:

$$\left[\frac{1}{G}\frac{dG}{dr_*}\right]_{r_*=R_*} = \frac{2(4n+1)M-(5n+2)R}{(n+2)R^2} - \frac{nR^2\omega^2}{(n+2)M}. \tag{B17}$$

Thus, a standard Sturm-Liouville eigenvalue system is established and the normalized eigenfunctions of (B13), $G_n(r_*)$, where $n = 1, 2, 3, \ldots$, form a complete set and satisfy the orthogonality relation (see, e.g., [43]):

$$\int_0^{R_*} \frac{p^2 G_m(r_*)G_n(r_*)}{e^{\nu}(\rho+p)C_s^2} dr_* = \delta_{mn}. \tag{B18}$$

4. Numerical results

The approximate real eigenfrequencies σ_n obtained from the CA scheme (B2) are listed and compared with the exact values $\omega_{\rm r}$ in Tables I, II and III. For p-modes of NSs constructed with realistic EOSs, e.g., A and SLy (see Tables I and II), the percentage error in $\omega_{\rm r}$ is usually less than a few percent and decreases with the mode order and the compactness of the star. However, the accuracy of the CA scheme (B2) could substantially worsen for polytropic stars (especially those with large polytropic indices) in spite of the fact that it still improves with increasing compactness and mode order (see Table III). In comparison with the two CA schemes proposed in the *H-S* approach, i.e., (7.2) and (7.1), (B2) is in general the least accurate one.

Moreover, it is worthwhile to note that the CA scheme (B2) fails to locate the f-mode. Hence, the n-th eigenfunction G_n $(n = 1, 2, 3, \ldots)$ in fact corresponds to the n-th p-mode. In Fig. 3 we show the scaled eigenfunction $\tilde{G}_n$ defined as:

$$\tilde{G}_n(r_*) \equiv \frac{pG_n(r_*)}{\sqrt{e^{\nu}(\rho+p)C_s^2}}, \tag{B19}$$

which satisfies the orthonormal condition $\int_0^{R_*} \tilde{G}_m(r_*)\tilde{G}_n(r_*)dr_* = \delta_{mn}$. There are $n-1$ nodes (excluding the two endpoints $r = 0$ and $r = R$) in the wave function $\tilde{G}_n$ or equivalently G_n. In particular, the wave function of the p_1-mode is nodeless. This observation further confirms that the p_1-mode is indeed the ground state of the present eigen-system (see, e.g., [43]) and explains why the f-mode is missing in the CA scheme (B2) of the *F-S* approach.

[1] N. K. Glendenning, *Compact Stars - Nuclear Physics, Particle Physics, and General Relativity* (Springer, NY, 1997).
[2] J. Lattimer and M. Prakash, Phys. Rep. **442**, 109 (2007).
[3] F. Weber, *Pulsars as Astrophysical Laboratories for Nuclear and Particle Physics* (Institute of Physics Publishing, 1999).
[4] N. Andersson and K. D. Kokkotas, Phys. Rev. Lett **77**, 4134 (1996).
[5] N. Andersson and K. D. Kokkotas, Mon. Not. R. Astron. Soc. **299**, 1059 (1998).
[6] O. Benhar, E. Berti, and V. Ferrari, Mon. Not. R. Astron. Soc. **310**, 797 (1999).
[7] K. D. Kokkotas, T. A. Apostolatos, and N. Andersson, Mon. Not. R. Astron. Soc. **320**, 307 (2001).
[8] O. Benhar, V. Ferrari, and L. Gualtieri, Phys. Rev. D **70**, 124015 (2004).
[9] L. K. Tsui and P. T. Leung, Phys. Rev. Lett. **95**, 151101 (2005).
[10] L. K. Tsui, P. T. Leung, and J. Wu, Phys. Rev. D **74**, 124025 (2006).
[11] H. K. Lau, P. T. Leung, and L. M. Lin, Astrophys. J. **714**, 1234 (2010).
[12] H. Sotani, N. Yasutake, T. Maruyama, and T. Tatsumi, Phys. Rev. D **83**, 024014 (2011).
[13] S. Hughes, Ann. Phys. **303**, 142 (2003).
[14] C. L. Fryer, D. E. Holz, and S. A. Hughes, Astrophys. J. **565**, 430 (2002).
[15] N. Andersson *et al.*, Gen. Rel. Grav. **43**, 409 (2011).
[16] S. Hild *et al.*, Class. Quant. Grav. **28**, 094013 (2011).
[17] K. S. Thorne and A. Campolattaro, Astrophys. J. **149**, 591 (1967).
[18] L. Lindblom and S. L. Detweiler, Astrophys. J. **53**, 73 (1983).
[19] S. Detweiler and L. Lindblom, Astrophys. J. **292**, 12 (1985).
[20] S. Chandrasekhar and V. Ferrari, Proc. R. Soc. Lond. A **432**, 247 (1991).
[21] L. Lindblom, G. Mendell, and J. R. Ipser, Phys. Rev. D. **56**, 2118 (1997).
[22] K. D. Kokkotas and B. G. Schmidt, Living Rev. Rel. **2**, 2 (1999).
[23] G. Allen, N. Andersson, K. D. Kokkotas, and B. F. Schutz, Phys. Rev. D **58**, 124012 (1998).
[24] W. Press, Astrophys. J. **170**, L105 (1971).
[25] E. W. Leaver, Phys. Rev. D **34**, 384 (1986).
[26] E. S. C. Ching, P. T. Leung, A. M. van den Brink, W. M. Suen, S. S. Tong, and K. Young, Rev. Mod. Phys. **70**, 1545 (1998).
[27] K. D. Kokkotas and B. G. Schmidt, Living Rev. Rel. **2**, 2 (1999).
[28] E. Berti, V. Cardoso, and A. O. Starinets, Class. Quant. Grav. **26**, 163001 (2009).
[29] T. G. Cowling, Mon. Not. R. Astron. Soc. **101**, 367 (1941).
[30] L. Lindblom and R. J. Splinter, Astrophys. J. **348**, 198 (1990).
[31] N. Andersson, K. D. Kokkotas, and B. F. Schutz, Mon. Not. R. Astron. Soc. **280**, 1230 (1996).
[32] J. Wu and P. T. Leung, Mon. Not. R. Astron. Soc. **381**, 151 (2007).
[33] P. M. Morse and H. Feshbach, *Methods of Theoretical Physics, Part I* (McGraw-Hill, 1953).
[34] J. R. Oppenheimer and G. M. Volkoff, Phys. Rev. **55**, 374 (1939).
[35] R. C. Tolman, Phys. Rev. **55**, 364 (1939).
[36] T. Regge and J. A. Wheeler, Phys. Rev. **108**, 1063 (1957).
[37] R. Price and K. S. Thorne, Astrophys. J. **155**, 163 (1969).
[38] F. J. Zerilli, Phys. Rev. Lett. **24**, 737 (1970).
[39] G. Allen, N. Andersson, K. D. Kokkotas, and B. F. Schutz, Phys. Rev. D **58**, 124012 (1998).
[40] J. Ruoff, Phys. Rev. D **63**, 064018 (2001).
[41] K. S. Thorne, Astrophys. J. **158**, 997 (1969).
[42] V. Pandharipande, Nucl. Phys. A **174**, 641 (1971).
[43] A. Zettl, *Sturm-Liouville Theory*, Mathematical surveys and monographs (American Math. Soc., 2005).
[44] F. Douchin and P. Haensel, Astron. Astrophys. **380**, 151 (2001).
[45] P. Haensel and A. Y. Potekhin, Astron. Astrophy. **428**, 191 (2004).
[46] Y. J. Zhang, J. Wu, and P. T. Leung, Phys. Rev. D **83**, 064012 (2011).

TABLE I: Exact QNM frequency $(\omega_{\rm r}, \omega_{\rm i})$ and the CA frequencies σ_n obtained respectively from (7.2) and Ref. [21] are compared for the leading fluid modes of a EOS A star ($\mathcal{C} = 0.27$).

Mode	Exact	(7.2)	(7.1)	(B2)	Ref. [21]
	$(\omega_{\rm r},\, \omega_{\rm i})$	σ_n	σ_n	σ_n	σ_n
f	(0.1447, 7.541E-5)	0.1163	NA	NA	0.1665
p_1	(0.3932, 3.047E-6)	0.3928	0.3934	0.3903	0.3984
p_2	(0.6130, 5.824E-7)	0.6135	0.6130	0.6126	0.6159
p_3	(0.7639, 5.269E-8)	0.7640	0.7639	0.7636	0.7664

TABLE II: Exact complex eigenfrequencies $(\omega_{\rm r}, \omega_{\rm i})$ and approximate real eigenfrequencies σ_n obtained respectively from CA (7.2), (7.1) and (B2) are listed for the leading fluid modes of NSs (EOS: Sly [44, 45]) with different compactness $\mathcal{C}$. Entries labelled by 'NA' signify modes that cannot be located by a specific approximation.

$\mathcal{C}$	Mode	Exact	(7.2)	(7.1)	(B2)
		$(\omega_{\rm r}, \omega_{\rm i})$	σ_n	σ_n	σ_n
0.117	f	(0.0488, 1.162E-5)	0.0410	NA	NA
0.117	p_1	(0.1654, 4.866E-7)	0.1651	0.1654	0.1628
0.117	p_2	(0.2079, 2.029E-8)	0.2079	0.2079	0.2071
0.117	p_3	(0.2390, 5.709E-9)	0.2390	0.2390	0.2383
0.210	f	(0.1047, 5.290E-5)	0.0848	NA	NA
0.210	p_1	(0.3281, 2.371E-6)	0.3279	0.3283	0.3245
0.210	p_2	(0.4995, 3.435E-8)	0.4996	0.4995	0.4985
0.210	p_3	(0.6154, 6.028E-9)	0.6155	0.6154	0.6150
0.276	f	(0.1476, 7.570E-5)	0.1177	NA	NA
0.276	p_1	(0.4074, 1.884E-6)	0.4072	0.4075	0.4053
0.276	p_2	(0.6236, 6.806E-7)	0.6241	0.6236	0.6234
0.276	p_3	(0.8192, 7.455E-8)	0.8195	0.8192	0.8193

TABLE III: Exact complex eigenfrequencies $(\omega_{\rm r}, \omega_{\rm i})$ and approximate real eigenfrequencies σ_n obtained respectively from CA (7.2), (7.1) and (B2) are listed for the leading fluid modes of polytropic stars (EOS: $p = 100\rho^{1.8}$) with different compactness $\mathcal{C}$. Entries labelled by 'NA' signify modes that cannot be located by a specific approximation.

$\mathcal{C}$	Mode	Exact	(7.2)	(7.1)	(B2)
		$(\omega_{\rm r}, \omega_{\rm i})$	σ_n	σ_n	σ_n
0.117	f	(0.0525, 1.329E-5)	0.0475	NA	NA
0.117	p_1	(0.1160, 3.283E-6)	0.1149	0.1164	0.1074
0.117	p_2	(0.1761, 2.680E-7)	0.1757	0.1762	0.1727
0.117	p_3	(0.2338, 1.473E-8)	0.2336	0.2338	0.2320
0.204	f	(0.1160, 5.771E-5)	0.1021	NA	NA
0.204	p_1	(0.2337, 1.728E-5)	0.2307	0.2343	0.2202
0.204	p_2	(0.3477, 5.490E-7)	0.3470	0.3478	0.3429
0.204	p_3	(0.4584, 3.069E-8)	0.4582	0.4584	0.4561
0.271	f	(0.1659, 5.795E-5)	0.1473	NA	NA
0.271	p_1	(0.3012, 1.341E-5)	0.2948	0.3020	0.2925
0.271	p_2	(0.4249, 2.832E-5)	0.4242	0.4250	0.4249
0.271	p_3	(0.5496, 1.359E-5)	0.5498	0.5495	0.5501

TABLE IV: QNM frequencies obtained from various ICA schemes for the leading polar w-modes of a EOS A star ($\mathcal{C} = 0.27$). The first two modes are classified as the $w_{\rm II}$ modes. Entries labelled by 'NA' signify modes that cannot be located by a specific approximation.

Exact	Exact	ICA	ICA	HFICA	HFICA	Ref. [21]	Ref. [21]
$\omega_{\rm r}$	$\omega_{\rm i}$	$\omega_{\rm r}$	$\omega_{\rm i}$	$\omega_{\rm r}$	$\omega_{\rm i}$	$\omega_{\rm r}$	$\omega_{\rm i}$
0.2067	0.7935	0.2065	0.7932	0.3304	0.8922	NA	NA
0.4593	0.3891	0.4599	0.3884	0.5076	0.4980	NA	NA
0.5096	0.1777	0.5095	0.1781	0.5102	0.1131	0.4348	0.0605
0.8730	0.3025	0.8730	0.3028	0.8335	0.2781	0.7924	0.1244
1.2266	0.3507	1.2266	0.3508	1.2006	0.3416	1.1446	0.1555
1.5782	0.3837	1.5782	0.3838	1.5589	0.3788	1.4961	0.1749
1.9302	0.4093	1.9301	0.4093	1.9148	0.4062	1.8476	0.1892
2.2828	0.4308	2.2828	0.4308	2.2700	0.4287	2.2004	0.2013
2.6359	0.4500	2.6359	0.4500	2.6250	0.4485	2.5537	0.2122
2.9894	0.4682	2.9894	0.4683	2.9799	0.4670	2.9079	0.2231
3.3429	0.4860	3.3428	0.4860	3.3344	0.4850	3.2619	0.2347
3.6960	0.5038	3.6960	0.5038	3.6884	0.5030	3.6158	0.2468



(a)

(b)

(c)

FIG. 1: The normalized H-functions for (a) the f-mode; (b) the p_1-mode; and (c) the p_2-mode of a star constructed with EOS A ($\rho_c = 2.227 \times 10^{-3}\ \mathtt{km}^{-2}$, $\mathcal{C} = 0.27$) are plotted against the normalized radius r/R. The solid and the dashed lines are respectively the real and the imaginary parts of H-function under CA (7.2). The dots and crosses respectively show the real and the imaginary parts of the exact H-function.



FIG. 2: The function $D(r, \omega)$ is plotted against r/R with $\omega = \sigma_0, \sigma_1, \sigma_2$.



FIG. 3: The scaled function $\tilde{G}_n$ of a NS with SLy EOS [44, 45] and compactness $\mathcal{C} = 0.276$ is plotted against r/R for $n = 1$ (solid line), 2 (dashed line), 3 (dot-dashed line), which correspond to p_1, p_2 and p_3 modes, respectively.